

No 70
Stn 246 Ave St.
Inaugural Essay

on
Scarlatina

Read March 1828

For the degree of Doctor of Medicine
in the

University of Pennsylvania
by

Hudson S. Burr

New Jersey

January 20th 1828.

Scarlatina or Scarlet Fever

Scarlet fever is probably a disease of very modern origin. No mention of it is made by the ancient or Arabian authors, and the first time it is distinctly noticed is but little more than two hundred years ago. It has been suspected that the contagion came originally from Africa.

Be this as it may, it first broke out in a severe form in Spain in 1610, from whence it spread to Naples, where it raged epidemically in 1618. In 1689 the same disease made its appearance in London, and was described by Doctor Morton, though not with the accuracy of the first Spanish and Italian authors.

In 1753 it broke out in North America, and spread gradually but slowly over that continent. One of the most curious circumstances in the history of the disease is the slowness of its diffusion.

When the scarlet fever first appeared in Europe, it was in a very malignant form; but between the years 1660

and 1670, a febrile disease attended with scarlet eruption was observed by Sydenham in a form singularly mild, that nosologist have doubted its being really the same disease with that which had previously occurred.

Dr Cullen believed it was specifically different.

Dr Withering states, that in his early practice he considered scarlet fever and putrid sore throat distinct diseases, requiring distinct methods of treatment. More enlarged experience however compelled him to renounce that opinion; and he says, that after paying the assiduous attention to the subject, by observing the disease in every difference of season, exposure, age and temperament, he was satisfied that they constitute but one species of disease;—that the variations in their appearance depend upon contingent circumstances, and that their greatest differences are not greater than those of the distinct and confluent small-pox.² ("Googery.") I have not the least doubt but it was a translation of the simple of our present nomenclature.

[Faint, illegible handwriting on the left page]

Charles
for the
company
It is m
then in
attacked
Charles
14. J
2nd J
3rd J
is a calc
and Patre
Charles
tional de
Charles
of the san
and infl
Charles
of the san

Scarlatina receives its name from the peculiarly florid appearance of the skin by which it is accompanied.

It is more prevalent in winter and in spring, than in summer and autumn, and more frequently attacks children than adults.

Scarlatina is divided into three forms viz:

1st. Scarlatina Simplex

2nd. Scarlatina Anginosa

3rd. Scarlatina Maligna; this third form is also called Cynancha Maligna or Putrida and Putrid Sore Throat.

Scarlatina Simplex is the simple constitutional disease.

Scarlatina Anginosa is a higher degree of the same disease, the throat being swollen and inflamed.

Scarlatina Maligna is a still higher degree of the same disease, being of a malignant character.

[Faint, mirrored handwriting from the reverse side of the page, appearing as bleed-through.]

the Royal
These are
They are re
consequent
Gravities a
Qualities
and ter
mastic an
wise accor
paid the
red there
paid in t
of the inter
slight the
over sever
Reliqua
the
patient
that facie

The throat is inflamed in this species also.

These are the same disease in different degrees, they are related to each other as the distinct and confluent small pox, typhus mitis and typhus gravior and the several stages of bilious fever.

Scarlatina comes on with anorexia, shivering, ful-
min and tension of the head, prostration of strength,
nausea and vomiting, the skin is hot, fluid or others-
wise according to the nature of the case. At this
period the throat is slightly inflamed, the tongue
red, there is more or less of a catarrhal affection;
pain in the head, and sometimes a disturbance
of the intellectual faculties. The disease being
slight the complaint will be "Simplex" when
more severe "Anginosa" and when threatening
"Maligna".

In the malignant form of the disease, the
patient becomes pale, sick and faint, the intelle-
tual faculties confused, praecordial oppression,

pa pitale
sa corre
contents
m die p
spots of
exhibit a
her lakes
on thum
Such s
Portiguo
to some
m dng
a harae
the disco
and sour
the form
the cry
patches
st appe

palpitation of heart, uneasiness of stomach without vomiting, its power being prostrated, its contents are however thrown up by a spasmodic singultus. The surface is pale with spots of a livid or dark mahogany hue, the eyes exhibit a fuliginous or insubriated expression—death soon takes place, the system being overwhelmed or stunned by a shock it cannot resist.

Such is the progress of a fatal case of Scarlatina Maligna; but in ordinary cases the skin begins to become red, with an efflorescence and there is swelling of the fauces, the pulse then exhibits a character belonging to the commencement of the disease, —the temperature of the body, thirst and scurpen the tongue graduated according to the form of the disease.

The eruption assumes the form of red points or patches which spread over the adjacent part. It appears first on the head and neck, then

It is ad
of the de
Diagn
four is
but may
ception
tion of the
character
the dwe
In mea
the it is
numer
interm
for my
cooling
atlet of
In the pu
dth part
is and a

It is admitted that the susceptibility to attacks of the disease is diminished.

Diagnosis. The only disease with which scarlet fever is liable to be confounded is measles. From this it may be distinguished by the character of the eruptive fever, the colour of the skin, and the affection of the fauces.

Character of the eruptive fevers. The efflorescence of this disease appears generally on the second day. In measles; seldom till the fourth. In the scarlatina it is more full, spreads more and consists of innumerable points and spots under the cuticle intermixed with minute pimples, in some places forming continuous irregular patches, in others coalescing with a uniform flash over a considerable extent of surface.

In the measles the rash is composed of circular dots, partly distinct, partly set in small clusters or patches and a little elevated so as to give the sensation of

roughness
patches as
of rescent
circle w
2. Colour of
in the face
in the pier
2. Affection
is short, w
dilate,
of the tongue
paled of
4. Affection
of the eyes
theiliary
the apper
in muscle
motion of
run tears, a

roughness on passing the finger over them; These patches are seldom confluent, but form a number of crescents with large intervening portions of cuticle which retain their usual appearance.

2. Colour of the efflorescence. The colour of the rash in the scarlet fever is like that of a bilious lobster shell; in the measles a dark reddish brown like a raspberry.

3. Affection of the fauces. In scarlatina the cough is short, without expectoration; In measles, it is obstinate, harsh, and attended with the discharge of a tough acrimonious phlegm, coming up in repeated efforts.

4. Affection of the eyes. In scarlatina, the redness of the eyes is not attended with intolerance of light, the ciliary glands are not affected; and though they appear watery and shining, they never overflow. In measles, the insensibility to light, and the inflammation of the eyes and eyelids is considerable. The eyes run tears, and there is great oedema, and sneezing.

St. Charles
my attend
time, except
The pursuit
good dig
the course
lady floor
bar cases
Decon
dis. ³ St. Charles
pauance g
paired co
imperfect
come to a
day error
ple. wh
Where the
on the this
for force o

In scarlatina, depression, anxiety and great weakness attends the fever; in measles, inflammatory symptoms, except in the typhoid variety.

The peculiarity of the tongue in scarlatina is a good diagnostic, it is heavily increased, and through this crust the papillae shoot up, and appear peculiarly florid, they are considerably lengthened in bad cases. —

Prognosis. Great prostration of strength, bima, distillation of acrid sanies from the fauces, purple appearance of the fauces, with dark sloughs, and accompanied with diarrhoea, permanent recession, or an imperfection of the eruptions, a change of the fluorescence to a mahogany colour are fatal or exceedingly dangerous symptoms. In some cases death has taken place when there had been every appearance of recovery. Where the case is very malignant death will occur on the third or fourth day, but sometimes it lingers for three or four weeks, but generally it may be said

[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]

that the p
The signa
in a, the a
the skin, a
of the tany
fading off
case sleep a
Dissec
no and h
then each is
the visce
lots in its
When the
and seen, a
whole all
Bethole
this is inde
ting, the
many of the

that the patient is safe, if he passes the ninth day. The signs of a favourable termination are lubricious urine, the abatement of the heat and fiery manes of the skin, desquamation of the cuticle and the subsidence of the tumefactions, greater slowness of the pulse; the falling off of the sloughs; filling up of the ulcerations, easy sleep and return of appetite.

Dissections. The mucous membrane of the primæ viæ and lungs is found much phlogosed, that of the stomach is sometimes gangrenous.—congestions of the viscera are also observed, the brain is implicated both in its meninges and in its substance.

When the throat has been affected patches of gangrene are seen, and which sometimes extend through the whole alimentary canal to the anus.

Pathology. Scarlatina is of gastric origin: this is indicated by the anorexia, nausea and vomiting, the appearance on digestion confirms this view of the pathology of the disease. The other symptoms

as all
Treaty
calls for
and or
colonel
not diag
the Amer
of the alle
Tatar En
called for
taken in
not been
the names
tion that
against the
has live
The inst
cal mel,
cal. latio

are all secondary.

Treatment. In its simple form scarlatina calls for little medical aid: An Emetic of Ipecacuanha or Tartar Emetic should be given: a purge of calomel followed by the neutral salts, and use the mild diaphoretics. In the Anginose form the treatment is more vigorous, attend to the evacuation of the alimentary canal, active vomiting with Tartar Emetic should be instituted, unless bleeding is called for. The Emetic practice is highly efficacious, taken in the commencement, where the system has not been brought into a diseased state, is effectual in the remedy in breaking up the train of morbid association that the attack of the disease offers no security against the recurrence of it, hence some physicians have hesitated in giving emetics.

The emetic should be followed by purging with calomel, this is the most proper purge as it is best calculated to prevent congestions of the abdominal

and era in
bowels have
laxatives to
the bowels
and hard
appearance
the la dose
The warm
and I have
in the early
freezing the
to, is affords
dysentery
spasms
in the system
next to this
The warm
and has no
has been found

viscera which are very apt to occur. When the bowels have been fully evacuated, use the gentle laxatives to keep them in a soluble state.

If the bowels having been cleansed, if the pulse is full and hard resort to V.S. the relief afforded by it the appearance of the blood, and its effects on the pulse should direct the practitioner as to its propriety.

The warm and cold bath have both been recommended. "I have used neither says professor Chapman" in the early and in inflammatory stages of the disease, sponging the body with cold water may be resorted to, it affords great relief, immersion in cold water is objectionable as in some cases it has proved fatal. Sponging with cold water when there is much energy in the system and heat of surface is preferable, and next to this dashing cold water on the body.

The warm bath is recommended where the effluence has moved or imperfectly come out, but what has been found very useful under such circumstances

[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]

is a little
low a note
But there
can be no
any further
than apper
pot. lues
a cloth or
ad by dis
interior
Corydalis
indicated by
mangaes
alterative
It's chance
Serice, the
diligent g
accumulated
New The p

is a blister to the Esophagus and superficial to the lower extremities.

But there are local affections which also require care: when the throat is affected volatile liniment.

Serpentine leeches and blisters are to be used.

Warm applications to the throat afford relief, as hot potatoes mashed and applied round the neck in a cloth or hot mush used in a similar way. They act by diverting the disease from the interior to the exterior.

Congestions of the abdominal viscera, which are indicated by a sense of weight and pain, are to be managed by topical bleeding, blisters, and the alterative use of mercury.

To cleanse the foul ulcers of the throat vomiting is of service, the emetic is to be succeeded by the use of some detergent gargle, as a decoction of bark with myrrh, acidulated barley water, or an infusion of cayenne pepper. The following is recommended.

U. S. Sen. of
the let for
moderate,
the article
for time, be
In Germany
as a power
of physics;
he died a
fection in
expressed
infected;
stronger po
all these ex
and who de
sublet the
tensions an
that if the
right or m

Rp. Cayenne pepper 2 table spoonfulls

Common salt - 1

Boiling water

Vinegar each 3ij.

Infuse and strain, Give a table spoonfull as often as is required. This is used as a gargle and given internally by the practitioners of the west Indies in Angina maligna.

The treatment of the malignant form is nearly the same: the primæ viæ are to be evacuated by an Emetic - and a purge of calomel is to be given: bleeding is hardly ever advisable. When sinking comes on support the system by Carbonate of Ammonia, Camphor, Spirit of Rosegerm, bark and wine, and the ordinary external applications.

Long after the cure of the disease there are certain sequelæ which may require attention: Deafness may occur arising from the Eustachian tube being inflamed and thickened - time removes this affection.

Sp. U.

Infuse and
ster as is
internally
in an jar
The treat
arous the p
matic - co
hardly co
support to
for Spec
ordinary e
long after
equala wh
ear and
famed and

Obdema of the lower extremities frequently follows scarlet fever; the best treatment for this is bleeding moderate purgings and the exhibition of Digitalis; this article will sometimes prove eminently beneficial; frictions, bandages and exercise are to be used.

¹⁴ In Germany, the use of the belladonna has been proposed as a preventive for this disease. Its author is Hahneman of Leipzig; Berndt of Custrin states that out of one hundred and ninety-five cases of children under fourteen who took belladonna, and were freely exposed to the contagion fourteen only were infected: and that when he afterwards used a stronger preparation of the drug, every one escaped: all those exposed in families, who had the scarlatina and who did not take the medicine took the disease: whilst those who did, escaped. Other strong testimonies are given in its favour. Kopp of Berlin states that if the belladonna be taken in proper doses for eight or nine days before exposure, the persons

taking it
There grow
firman
given to a
more for
effect is p
an expe
black m
and of la
the disea

* E. Med. a

taking it are safe. The quantities given are very small: three grains of the extract are dissolved in an ounce of cinnamon water, and of this two or three drops are given to a child under twelve months and one drop more for every year above that. In general no effect is produced by it: Sometimes however it produces an eruption like that of scarlatina: It renders the attack more mild, if it does not prevent the disease; and if taken four or five days before exposure the disease never proves fatal."^o

* Ed. Med. and Surg. Journal January 1825.

To the Honorable